

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
1(i)	<p>Assume that a and b are non-parallel vectors.</p> <p>$\overrightarrow{OC} = 5\mathbf{a}$, $\overrightarrow{OD} = 2\mathbf{b}$</p> <p>On the line BC, $\overrightarrow{OM} = \lambda(5\mathbf{a}) + (1 - \lambda)\mathbf{b}$</p> <p>On the line AD, $\overrightarrow{OM} = \mu(2\mathbf{b}) + (1 - \mu)\mathbf{a}$</p> <p>Since a and b are non-zero, non-parallel vectors, comparing coefficient</p> $5\lambda = 1 - \mu$ $2\mu = 1 - \lambda \Rightarrow \lambda = \frac{1}{9}, \mu = \frac{4}{9}$ <p>Thus $\overrightarrow{OM} = \frac{5}{9}\mathbf{a} + \frac{8}{9}\mathbf{b}$</p>
1(ii)	<p>Since a is a unit vector in the direction of OC,</p> <p>shortest distance $= \left \overrightarrow{OM} \times \mathbf{a} \right$</p> $= \left \left(\frac{5}{9}\mathbf{a} + \frac{8}{9}\mathbf{b} \right) \times \mathbf{a} \right $ $= \frac{8}{9} \mathbf{a} \times \mathbf{b} $ <p>$k = \frac{8}{9}$</p>
2(a)	<p>For sum to infinity to exist,</p> $ \tan \theta < 1$ $-1 < \tan \theta < 1$ $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$ $\frac{1}{1 - \tan \theta} > 2$ $0 < 1 - \tan \theta < \frac{1}{2}$ $\tan \theta > \frac{1}{2} \Rightarrow \theta > \tan^{-1} \frac{1}{2} \quad (\text{or } \theta > 0.464)$ <p>Since $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$,</p> <p>therefore $\{\theta \in \square \mid \tan^{-1} \frac{1}{2} < \theta < \frac{\pi}{4}\}$</p> <p>(or $\{\theta \in \square \mid 0.464 < \theta < 0.785\}$ or $\theta : (0.464, 0.785)$)</p>

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
2(b)	$u_1 = S_1 = 2 \Rightarrow a = 8$ $u_2 = S_2 - S_1 = 10 \Rightarrow d = 8$ $u_{32} = a + (32-1)d = 2 + (32-1)8 = 250$ $\frac{u_{32}}{u_m} = \frac{u_m}{u_2} = \text{constant}$ $\Rightarrow (u_m)^2 = (10)(250) = 2500$ $u_m = 50 \text{ (since it is a positive sequence)}$ $50 = 2 + (m-1)8 \Rightarrow m = 7$ <p>Alternatively,</p> $u_n = S_n - S_{n-1}$ $= 4n^2 - 2n - [4(n-1)^2 - 2(n-1)]$ $= 8n - 6$ $\frac{u_{32}}{u_m} = \frac{u_m}{u_2}$ $\frac{8(32) - 6}{8m - 6} = \frac{8m - 6}{8(2) - 6}$ $(8m - 6)^2 = (250)(10) = 2500$ $m = 7 \text{ or } m = -5.5 \text{ (rejected as } m \text{ is a positive integer)}$
3(i)	$y = \ln(\cos ax - \sin ax)$ $e^y = \cos ax - \sin ax$ $e^y \frac{dy}{dx} = -a \sin ax - a \cos ax$ $e^y \frac{d^2 y}{dx^2} + e^y \left(\frac{dy}{dx} \right)^2 = -a^2 \cos ax + a^2 \sin ax$ $e^y \frac{d^2 y}{dx^2} + e^y \left(\frac{dy}{dx} \right)^2 = -a^2 (\cos ax - \sin ax)$ $e^y \frac{d^2 y}{dx^2} + e^y \left(\frac{dy}{dx} \right)^2 = -a^2 e^y$ $\frac{d^2 y}{dx^2} + \left(\frac{dy}{dx} \right)^2 + a^2 = 0$

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
3(ii)	$\frac{d^3 y}{dx^3} + 2 \left(\frac{dy}{dx} \right) \frac{d^2 y}{dx^2} = 0$ <p>When $x = 0$, $y = 0$</p> $\frac{dy}{dx} = -a, \quad \frac{d^2 y}{dx^2} = -2a^2, \quad \frac{d^3 y}{dx^3} = -4a^3$ $y = -ax - a^2 x^2 - \frac{2}{3} a^3 x^3 + \dots$
3(iii)	$\ln(\cos 2x - \sin 2x) = -2x - 4x^2 - \frac{16}{3} x^3 + \dots$ $\cos 2x - \sin 2x \approx e^{-2x-4x^2}$ $\approx 1 + (-2x - 4x^2) + \frac{(-2x - 4x^2)^2}{2!} \text{ (since } e^x \approx 1 + x + \frac{x^2}{2!} \text{)}$ $\approx 1 - 2x - 4x^2 + \frac{(-2x)^2}{2}$ $= 1 - 2x - 2x^2 \quad \text{where } k = -2$
3(iv)	$\cos 2x - \sin 2x = 1 - \frac{(2x)^2}{2} - (2x)$ $= 1 - 2x - 2x^2$

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
4(i)	$\frac{dx}{dt} = k(N-x)x$ $\frac{1}{(N-x)x} \frac{dx}{dt} = k$ $\int \frac{1}{(N-x)x} dx = \int k dt$ $\frac{1}{N} \int \frac{1}{N-x} + \frac{1}{x} dx = \int k dt$ $\frac{1}{N} (-\ln N-x + \ln x) = kt + C$ $\frac{1}{N} \ln \left \frac{x}{N-x} \right = kt + C$ $\ln \left \frac{x}{N-x} \right = Nkt + NC$ $\left \frac{x}{N-x} \right = e^{Nkt+NC}$ $\frac{x}{N-x} = Ae^{Nkt} \quad \text{where } A = \pm e^{NC}$ <p>When $t=0$, $x = \frac{1}{3}N$,</p> $A = \frac{1}{2}$ $\frac{x}{N-x} = \frac{1}{2} e^{Nkt}$ $2x = (N-x)e^{Nkt}$ $x(2 + e^{Nkt}) = Ne^{Nkt}$ $x = \frac{Ne^{Nkt}}{2 + e^{Nkt}} \quad \text{equivalently, } x = \frac{N}{2e^{-Nkt} + 1}$
	<u>Alternative method</u> (not recommended):

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
	$\frac{1}{(N-x)x} \frac{dx}{dt} = k$ $-\int \frac{1}{x^2 - Nx} dx = \int k dt$ $-\int \frac{1}{\left(x - \frac{N}{2}\right)^2 - \left(\frac{N}{2}\right)^2} dx = \int k dt$ $-\left(\frac{1}{2\left(\frac{N}{2}\right)} \ln \left \frac{\left(x - \frac{N}{2}\right) - \frac{N}{2}}{\left(x - \frac{N}{2}\right) + \frac{N}{2}} \right \right) = kt + C$ $-\left(\frac{1}{N} \ln \left \frac{x - N}{x} \right \right) = kt + C$ $\frac{1}{N} \ln \left \frac{x}{N - x} \right = kt + C$ $\ln \left(\frac{x}{N - x} \right) = Nkt + NC \quad \text{since } 0 < x < N$ <p>When $t=0$, $x = \frac{1}{3}N$, $\ln \frac{1}{2} = NC \Rightarrow C = -\frac{1}{N} \ln 2$</p> $\ln \left(\frac{x}{N - x} \right) = Nkt - \ln 2$ $\ln \left(\frac{2x}{N - x} \right) = Nkt$ $\frac{2x}{N - x} = e^{Nkt}$ $x(2 + e^{Nkt}) = Ne^{Nkt}$ $x = \frac{Ne^{Nkt}}{2 + e^{Nkt}} \quad \text{equivalently, } x = \frac{N}{2e^{-Nkt} + 1}$ <p>As $t \rightarrow \infty$, $x = \frac{N}{2e^{-Nkt} + 1} \rightarrow N$</p>

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn													
(ii)	$\frac{d^2 x}{dt^2} = \frac{-9t}{(4+9t^2)^2}$ $\int \frac{d^2 x}{dt^2} dt = \int \frac{-9t}{(4+9t^2)^2} dt$ $= -\frac{1}{2} \int \frac{18t}{(4+9t^2)^2} dt$ $\frac{dx}{dt} = \frac{1}{2} \left(\frac{1}{4+9t^2} \right) + A$ $\int \frac{dx}{dt} dt = \frac{1}{2} \int \frac{1}{4+9t^2} dt + \int A dt$ $= \frac{1}{18} \int \frac{1}{\left(\frac{2}{3}\right)^2 + t^2} dt + \int A dt$ $= \frac{1}{18} \cdot \frac{1}{\frac{2}{3}} \tan^{-1} \left(\frac{t}{\frac{2}{3}} \right) + At + B$ $x = \frac{1}{12} \tan^{-1} \left(\frac{3t}{2} \right) + At + B$ <p>Since the population stabilises in the long run, as $t \rightarrow \infty$, $x \rightarrow$ finite value , $A = 0$</p> <p>When $t = 0$, $x = \frac{1}{3}N$, $B = \frac{N}{3}$</p> <p>Hence $x = \frac{1}{12} \tan^{-1} \left(\frac{3t}{2} \right) + \frac{N}{3}$</p> <p>When $t \rightarrow \infty$, $\tan^{-1} \left(\frac{3t}{2} \right) \rightarrow \frac{\pi}{2}$</p> <p>Hence $x \rightarrow \frac{\pi}{24} + \frac{N}{3}$.</p>												
5(i)	<p>Let Y be the payment for an individual. The probability table is as follows:</p> <table><tr><td>y</td><td>100</td><td>200</td><td>300</td><td>325</td><td>350</td></tr><tr><td>$P(Y = y)$</td><td>$\frac{1}{3}$</td><td>$\frac{4}{15}$</td><td>$\frac{1}{5}$</td><td>$\frac{2}{15}$</td><td>$\frac{1}{15}$</td></tr></table> $E(Y) = 100 \cdot \frac{1}{3} + 200 \cdot \frac{4}{15} + 300 \cdot \frac{1}{5} + 325 \cdot \frac{2}{15} + 350 \cdot \frac{1}{15}$ $= 213\frac{1}{3}$	y	100	200	300	325	350	$P(Y = y)$	$\frac{1}{3}$	$\frac{4}{15}$	$\frac{1}{5}$	$\frac{2}{15}$	$\frac{1}{15}$
y	100	200	300	325	350								
$P(Y = y)$	$\frac{1}{3}$	$\frac{4}{15}$	$\frac{1}{5}$	$\frac{2}{15}$	$\frac{1}{15}$								


2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
5(ii)	$E(Y^2) = 100^2 \cdot \frac{1}{3} + 200^2 \cdot \frac{4}{15} + 300^2 \cdot \frac{1}{5} + 325^2 \cdot \frac{2}{15} + 350^2 \cdot \frac{1}{15}$ $= 54250$ $\text{Var}(Y) = E(Y^2) - [E(Y)]^2 = 8738\frac{8}{9}$ <p>Since $n=100$ is large, by Central Limit Theorem,</p> $T = \sum_{i=1}^{100} Y_i \sim N(21333.33, 873888.89) \text{ approx.}$ <p>Prob. Req'd = $P(T > 24000)$ ≈ 0.00217</p>
6	<p>Probability required = $\frac{2 \times {}^3C_2 \times {}^5C_2}{{}^8C_4} = \frac{6}{7}$</p>
6(i)	<p>Required probability = $\frac{(8-1)! {}^8C_2 2!}{(10-1)!} = \frac{7}{9}$</p> <p>OR $1 - \frac{(9-1)! 2!}{(10-1)!} = \frac{7}{9}$</p>
6(ii)	<p>Let X be the event that the remaining 3 girls are separated.</p> <p>Let Y be the event that Ann and Alice are not seated together.</p> $P(X Y) = \frac{P(X \cap Y)}{P(Y)}$ $= \frac{P(X) - P(X \cap Y')}{P(Y)}$ $= \frac{\frac{(7-1)! {}^7C_3 3! - (6-1)! 2! {}^6C_3 3!}{(10-1)!}}{\frac{7}{9}}$ $= \frac{\cancel{85} / \cancel{252}}{\cancel{7} / 9}$ $= \frac{85}{196}$

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn																																																																																																										
7(i)	$P(R=6)=2P(R=4)$ ${}^{10}C_6p^6(1-p)^4=2{}^{10}C_4p^4(1-p)^6$ $p^2=2(1-p)^2$ $p^2-4p+2=0$ $p=0.586$																																																																																																									
7(ii)	$R\sim B(n,0.25)$ $P(R<2)>0.15$ $P(R\leq1)>0.15$ $n=12$ <table><tr><th colspan="7">NORMAL FLOAT AUTO REAL RADIAN MP</th></tr><tr><th colspan="7">PRESS + FOR Δ Tb1</th></tr><tr><th>X</th><th>Y1</th><th></th><th></th><th></th><th></th><th></th></tr><tr><td>7</td><td>.44495</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>8</td><td>.36708</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>9</td><td>.30034</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>10</td><td>.24403</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>11</td><td>.1971</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>12</td><td>.15838</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>13</td><td>.12671</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>14</td><td>.10097</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>15</td><td>.08018</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>16</td><td>.06348</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>17</td><td>.05011</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td colspan="7">X=12</td></tr></table>	NORMAL FLOAT AUTO REAL RADIAN MP							PRESS + FOR Δ Tb1							X	Y1						7	.44495						8	.36708						9	.30034						10	.24403						11	.1971						12	.15838						13	.12671						14	.10097						15	.08018						16	.06348						17	.05011						X=12						
NORMAL FLOAT AUTO REAL RADIAN MP																																																																																																										
PRESS + FOR Δ Tb1																																																																																																										
X	Y1																																																																																																									
7	.44495																																																																																																									
8	.36708																																																																																																									
9	.30034																																																																																																									
10	.24403																																																																																																									
11	.1971																																																																																																									
12	.15838																																																																																																									
13	.12671																																																																																																									
14	.10097																																																																																																									
15	.08018																																																																																																									
16	.06348																																																																																																									
17	.05011																																																																																																									
X=12																																																																																																										
7(iii)	$R\sim B(11,0.7)$ $P(R\geq5 R\leq8)=\frac{P(R\geq5\text{ and }R\leq8)}{P(R\leq8)}$ $=\frac{P(5\leq R\leq8)}{P(R\leq8)}$ $=\frac{P(R\leq8)-P(R\leq4)}{P(R\leq8)}$ $=0.969$																																																																																																									
8(a)(i)	The value of 0.073 indicates that there is a weak linear correlation between petal size and the amount of water but there could be some non-linear relation.																																																																																																									
8(a)(ii)	The approximate value of -1 indicates that there is a strong negative linear correlation between the risk of heart disease and amount of red wine intake. It does not mean that red wine intake decreases the risk of heart disease.																																																																																																									

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
8(b) (i)	
8(b) (ii)	<p>For Model A, $r = -0.9985438 = 0.99854$</p> <p>For Model B, $r = 0.9984431 = 0.99844$</p>
8(b) (iii)	<p>Model A as the r value is closer to 1.</p> <p>The suitable regression line is $y = 848.24 - \frac{1629165.57}{x}$</p>
8(b) (iv)	<p>This is because age (x) is the controlled variable</p>
8(b) (v)	<p>The rise in the median age is due to the drop in the growth of the population.</p>
9(i)	<p>To test $H_0 : \mu = 0.5$ $H_1 : \mu \neq 0.5$ Level of significance: 5% Under H_0, $Z = \frac{\bar{X} - 0.5}{0.04 / \sqrt{25}} \sim N(0,1)$ Reject H_0 if p-value ≤ 0.05 Calculation: $\bar{x} = 0.51$, p-value = 0.211 Since p-value > 0.05, we do not reject H_0. Thus there is insufficient evidence at 5% level of significance that the manufacturing process is producing ball bearings of different diameters. Distribution of the diameter of the ball bearings is normal.</p>

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
9(ii)	<p>To test $H_0 : \mu = 0.5$ $H_1 : \mu > 0.5$ Level of significance: 5%</p> <p>Under H_0, by Central Limit Theorem, $Z = \frac{\bar{X} - 0.5}{s/\sqrt{100}} \sim N(0,1)$ approx</p> <p>Reject H_0 if p-value ≤ 0.05</p> <p>Calculation: $\bar{x} = 0.506$, $\Sigma(x - 0.5) = 50.6 - 50 = 0.6$</p> $s^2 = \frac{1}{n-1} \left[\Sigma(x-0.5)^2 - \frac{(\Sigma(x-0.5))^2}{n} \right]$ $= \frac{1}{99} \left(0.08345 - \frac{0.6^2}{100} \right)$ $= 8.07 \times 10^{-4}$ p -value = 0.0173 <p>Since p-value < 0.05, we reject H_0. Thus there is sufficient evidence at 5% level of significance that the manufacturing process is producing oversized ball bearings.</p>
9(iii)	<p>For the new sample, $\bar{x} = 0.506$. However,</p> $s^2 = \frac{1}{199} \left(2(0.08345) - \frac{(2(0.6))^2}{200} \right)$ $= 8.02513 \times 10^{-4}$ $Z_{calc} = 2.995$, p -value = 0.00137 <p>Since p-value < 0.05, we will still reject H_0. The conclusion remains the same.</p>
10(i)	<p>Let X denotes the diameter of bolt from manufacturer A. $X \sim N(1.56, 0.16^2)$</p> <p>Let Y denotes the diameter of bolt from manufacturer B. $Y \sim N(\mu, 0.16^2)$</p> <p>$P(Y < 1.52) = 0.242$</p> $P\left(Z < \frac{1.52 - \mu}{0.16}\right) = 0.242$ $\frac{1.52 - \mu}{0.16} = -0.6998836 \Rightarrow \mu = 1.63198 = 1.632$
10(ii)	<p>$W = X - Y \sim N(1.56 - 1.632, 0.16^2 + 0.16^2)$</p> <p>$P(W < 0.1) = P(-0.1 < W < 0.1) = 0.326$</p>

2017 NYJC JC2 Preliminary Examination 9758/2 Solution

Qn	
10(iii)	$X_1 + X_2 + X_3 + X_4 + X_5 - 5Y \sim N(5(1.56) - 5(1.632), 5(0.16^2) + 5^2(0.16)^2)$ $P(X_1 + X_2 + X_3 + X_4 + X_5 > 5Y)$ $= P(X_1 + X_2 + X_3 + X_4 + X_5 - 5Y > 0)$ $= 0.341$
10(iv)	$P(X < 1.52) = 0.40129$ $\text{Prob. Req'd} = (0.44)(0.4012) + 0.56(0.242)$ $= 0.3120876$ $= 0.312$