

## Section A

1

$$f(0) = \cos 0 - 0 = 1 > 0$$

$$f(1) = \cos 1 - 1 = -0.459 < 0$$

By Linear Interpolation, there is at least 1 root in the interval  $(0, 1)$ .

$$f'(x) = -\sin x - 1 < 0 \text{ in the interval } (0, 1).$$

$\therefore f(x)$  is strictly decreasing in the interval  $(0, 1)$ .

Hence there is only one root in the interval  $(0, 1)$ .

Given  $x_{n+1} = \cos x_n$ , and  $x_1 = 1$ ,

$$x_2 = 0.5403$$

$$x_3 = 0.8575$$

$$x_4 = 0.6542$$

$$x_5 = 0.7934$$

$$x_6 = 0.7013$$

$$x_7 = 0.7639$$

$$x_8 = 0.7221$$

$$x_9 = 0.7504$$

$$x_{10} = 0.7314$$

$$x_{11} = 0.7442$$

$$x_{12} = 0.7356$$

$$f(0.745) = -0.0099 < 0$$

$$f(0.735) = 0.0068 > 0$$

Hence,  $0.735 < \alpha < 0.745$

$$\therefore \alpha = 0.74$$

$$\begin{aligned}\tan \theta &= \frac{y}{x} \\ &= \frac{a \sin t - 2a \sin t \cos t}{a \cos t - 2a \cos^2 t} \\ &= \tan t\end{aligned}$$

$$\therefore \theta = t$$

$$\begin{aligned}r^2 &= x^2 + y^2 \\ &= (a \cos \theta)^2 (1 - 2 \cos \theta)^2 + (a \sin \theta)^2 (1 - 2 \cos \theta)^2 \\ &= (1 - 2 \cos \theta)^2 \left[ (a \cos \theta)^2 + (a \sin \theta)^2 \right] \\ &= a^2 (1 - 2 \cos \theta)^2\end{aligned}$$

$$\therefore r = a(1 - 2 \cos \theta)$$

When  $\theta = 0$ ,  $r = -a$ .

When  $\theta = \frac{\pi}{2}$ ,  $r = a$ .

When  $\theta = \pi$ ,  $r = 3a$ .

When  $r = 0$ ,  $\theta = \frac{\pi}{3}, -\frac{\pi}{3}$ .

$$\text{Area} = 2 \left[ \left| \int_{\frac{\pi}{3}}^{\pi} \frac{1}{2} r^2 d\theta \right| - \left| \int_0^{\frac{\pi}{3}} \frac{1}{2} r^2 d\theta \right| \right]$$

Note that  $\int_{\frac{\pi}{3}}^{\pi} \frac{1}{2} r^2 d\theta > 0$  and  $\int_0^{\frac{\pi}{3}} \frac{1}{2} r^2 d\theta < 0$

$$\begin{aligned}\text{Hence area} &= 2 \left[ \int_{\frac{\pi}{3}}^{\pi} \frac{1}{2} r^2 d\theta + \int_0^{\frac{\pi}{3}} \frac{1}{2} r^2 d\theta \right] \\ &= 2 \int_0^{\pi} \frac{1}{2} r^2 d\theta \\ &= a^2 \int_0^{\pi} (1 - 2 \cos \theta)^2 d\theta \\ &= a^2 \int_0^{\pi} 1 - 4 \cos \theta + 4 \cos^2 \theta d\theta \\ &= a^2 \int_0^{\pi} 1 - 4 \cos \theta + 4 \left( \frac{1 + \cos 2\theta}{2} \right) d\theta \\ &= a^2 \int_0^{\pi} 3 - 4 \cos \theta + 2 \cos 2\theta d\theta \\ &= a^2 [3\theta - 4 \sin \theta + \sin 2\theta]_0^{\pi} \\ &= 3\pi a^2\end{aligned}$$

<p>3(a) (i)</p>	<p>Using Euler's Method, we have</p> $y_{n+1} = y_n + 0.5 \left( e^{\sin t_n} + e^{\cos y_n} \right)$ <p>Given <math>t_0 = 0, y_0 = 0</math></p> $t_1 = 0.5$ $y_1 = 0 + 0.5 \left( e^{\sin 0} + e^{\cos 0} \right) = 1.85914$ $t_2 = 1$ $y_2 = 1.85914 + 0.5 \left( e^{\sin(0.5)} + e^{\cos(1.85914)} \right)$ $= 3.04295$ $= 3.043$
<p>3(a) (ii)</p>	<p>Using the improved Euler's Method</p> $y_{n+1}^* = y_n + 0.5 \left( e^{\sin t_n} + e^{\cos y_n} \right)$ $y_{n+1} = y_n + 0.5 \left[ \frac{\left( e^{\sin t_n} + e^{\cos y_n} \right) + \left( e^{\sin t_{n+1}} + e^{\cos y_{n+1}^*} \right)}{2} \right]$ <p>Given <math>t_0 = 0, y_0 = 0</math></p> $t_1 = 0.5$ $y_1^* = 0 + 0.5 \left( e^{\sin 0} + e^{\cos 0} \right) = 1.85914$ $y_1 = 0 + 0.5 \left[ \frac{\left( e^{\sin 0} + e^{\cos 0} \right) + \left( e^{\sin(0.5)} + e^{\cos(1.85914)} \right)}{2} \right] = 1.52148$ $y_2^* = 1.52148 + 0.5 \left( e^{\sin(0.5)} + e^{\cos(1.52148)} \right) = 2.85431$ $y_2 = 1.52148 + 0.5 \left[ \frac{\left( e^{\sin(0.5)} + e^{\cos(1.85914)} \right) + \left( e^{\sin(1)} + e^{\cos(2.85431)} \right)}{2} \right]$ $= 2.86366$ $= 2.864$

3(b)	<p>Given <math>h = \frac{2-0}{5-1} = \frac{1}{2}</math></p> <p>When <math>t = 0</math>, <math>v = e^{\sin 0} = 1</math></p> <p><math>t = 0.5</math>, <math>v = e^{\sin(0.5)} = 1.61514</math></p> <p><math>t = 1</math>, <math>v = e^{\sin(1)} = 2.31977</math></p> <p><math>t = 1.5</math>, <math>v = e^{\sin(1.5)} = 2.71148</math></p> <p><math>t = 2</math>, <math>v = e^{\sin(2)} = 2.48257</math></p> <p>Distanced Travelled <math>= \int_0^2 v \, dt</math></p> $\approx \frac{0.5}{3} [1 + 4(1.61514) + 2(2.31977) + 4(2.71148) + 2.48257]$ $= 4.238$
------	---

4(i)	$b^2 = a^2(1 - e^2)$ $4 = a^2(1 - 0.85^2)$ $a^2 = 14.4144$ $a = 3.7966$ $p = 2a = 7.59$
4(ii)	$\frac{PF}{Pd} = e$ $\frac{r}{r \cos \theta + d} = e$ $r = er \cos \theta + ed$ $r = \frac{ed}{1 - e \cos \theta} = \frac{k}{1 - 0.85 \cos \theta}$ <p>Sub <math>(0, a + ae)</math>.</p> $a + ae = \frac{k}{1 - 0.85}$ $k = (1 - 0.85)(3.796631)(1.85)$ $= 1.05303$
4(iii)	$\int_0^\pi \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta = 9.32060$ <p>Cost <math>= 9.32060 \times 6 \times 28 = \\$1566</math></p>

5(i)	$ M - \lambda I  = 0$ $(a - \lambda)^3 + 2(-2)^3 - 3(-2)^2(a - \lambda) = 0$ $(a - \lambda)^3 - 12(a - \lambda) - 16 = 0$ <p>Since <math>\lambda = -3</math> is a solution,</p> $(a + 3)^3 - 12(a + 3) - 16 = 0$ $a^3 + 9a^2 + 15a - 25 = 0$ <p>Using GC, <math>a = 1</math>, <math>a = -5</math>(rej)</p> $\therefore (1 - \lambda)^3 - 12(1 - \lambda) - 16 = 0$ $-\lambda^3 + 3\lambda^2 + 9\lambda - 27 = 0$ <p>Using GC, <math>\lambda = -3, 3, 3</math></p>
5(ii)	<p>When <math>\lambda = -3</math>,</p> $M - \lambda I = \begin{pmatrix} 4 & -2 & -2 \\ -2 & 4 & -2 \\ -2 & -2 & 4 \end{pmatrix}$ $\begin{pmatrix} 4 \\ -2 \\ 2 \end{pmatrix} \times \begin{pmatrix} -2 \\ 4 \\ -2 \end{pmatrix} = 2 \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}$ $v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$
5(iii)	<p>When <math>\lambda = -3</math>,</p> $M - \lambda I = \begin{pmatrix} -2 & -2 & -2 \\ -2 & -2 & -2 \\ -2 & -2 & -2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ $\tilde{x} = \begin{pmatrix} -s - t \\ s \\ t \end{pmatrix} = s \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$ $v_2 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \quad v_3 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$ <p> <math>l + m = n</math>  <math>l + m - n = 0</math>  <math>p + r = q</math>  <math>p - q + r = 0</math> </p>

	$Q = \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$
(iv)	$\begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$ $\left( \begin{array}{ccc c} 1 & -1 & -1 & 3 \\ 1 & 1 & 0 & 2 \\ 1 & 0 & 1 & 1 \end{array} \right) \rightarrow \left( \begin{array}{ccc c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right)$ $\therefore \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} - \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$ $\begin{aligned} M^k \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} &= M^k \left[ 2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} - \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right] \\ &= M^{k-1} \left( M \left[ \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} - \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right] \right) \\ &= M^{k-1} \left( 3 \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} - (-3) \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right) \\ &= M^{k-2} \left( 3^2 \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} - (-3)^2 \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right) \\ &\vdots \\ &= 3^k \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} - (-3)^k \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \\ &= 3^k \left[ \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} - (-1)^k \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right] \end{aligned}$

When  $k$  is even,

$$\begin{aligned} M^k \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} &= 3^k \left[ \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} - \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right] \\ &= 3^k \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} \end{aligned}$$

When  $k$  is odd,

$$\begin{aligned} M^k \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} &= 3^k \left[ \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} + \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right] \\ &= 3^k \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \end{aligned}$$

## Section B Marking Scheme

6(i)	<p>The two sample are not independent since they are taken on the same days. e.g. on public holidays, there is a larger crowd, so earnings for both restaurants will be higher.</p> <p>The assumptions are: <math>D_i</math>'s are independent and normally distributed.</p>																		
6(ii)	<p><math>H_0 : \mu_d = 0</math> <math>H_1 : \mu_d \neq 0</math> Test at <math>\alpha\%</math> sig level.</p> <p>Under <math>H_0</math>, test statistic is <math>T = \frac{\bar{D} - \mu_d}{s_d \sqrt{\frac{1}{12}}} \sim t_{11}</math></p> <p>Using GC, p-value = 0.0117</p> <p><math>H_0</math> rejected <math>\Rightarrow 0.0117 \leq \frac{\alpha}{100}</math></p> <p>Least <math>\alpha = 1.17</math></p>																		
7(i)	<p><math>H_0</math>: a person's colour of eyes and reaction to UV are independent. <math>H_1</math>: a person's colour of eyes and reaction to UV are not independent.</p> <p>Test at 5% sig level.</p> <p>Under <math>H_0</math>, expected freq is</p> <table><tr><td>15.675</td><td>7.425</td><td>9.9</td></tr><tr><td>26.125</td><td>12.375</td><td>16.5</td></tr><tr><td>15.2</td><td>7.2</td><td>9.6</td></tr></table> <p>Contributions to test statistic:</p> <table><tr><td>4.801</td><td>0.0445</td><td>6.6273</td></tr><tr><td>0.3164</td><td>0.456</td><td>0.015</td></tr><tr><td>2.2132</td><td>0.45</td><td>6.017</td></tr></table> <p>Using GC, <math>\chi^2 = 20.9</math> <math>p\text{-value} = 3.25 \times 10^{-4} &lt; 0.05</math> Reject <math>H_0</math> as there is sufficient evidence at 5% sig level that a person's eye colour and reaction to UV are not independent.</p> <p>The strongest evidence for association comes from the brown eye/ People with brown eyes tend to show no or just a slight reaction to UV light.</p> <p><i>diff b/w obs'd &amp; exp</i> <i>brown eye <math>\frac{O_i - E_i}{E_i}</math></i> <i>bigger the value the more evidence the factors not</i></p>	15.675	7.425	9.9	26.125	12.375	16.5	15.2	7.2	9.6	4.801	0.0445	6.6273	0.3164	0.456	0.015	2.2132	0.45	6.017
15.675	7.425	9.9																	
26.125	12.375	16.5																	
15.2	7.2	9.6																	
4.801	0.0445	6.6273																	
0.3164	0.456	0.015																	
2.2132	0.45	6.017																	
7(ii)	<p>sample proportion = 0.275 95% CI for <math>p = (0.1951, 0.3549)</math></p>																		

*diff btw obsd & expected.*

*brown eye  $\frac{O_i - E_i}{E_i^2}$*

*bigger the value the more evidence that two factors not indep.*



8(i)


$$\begin{aligned}
 P(A \leq a) &= P(\pi X^2 \leq a) \\
 &= P\left(X^2 \leq \frac{a}{\pi}\right) \\
 &= P\left(-\sqrt{\frac{a}{\pi}} \leq X \leq \sqrt{\frac{a}{\pi}}\right) \\
 &= P\left(1 \leq X \leq \sqrt{\frac{a}{\pi}}\right) \\
 &= \sqrt{\frac{a}{\pi}} - 1
 \end{aligned}$$

$$P(A = a) = \frac{d}{da} P(A \leq a)$$

$$\begin{aligned}
 &= \frac{1}{2} \left(\frac{a}{\pi}\right)^{-\frac{1}{2}} \frac{1}{\pi} \\
 &= \frac{1}{2\pi} \sqrt{\frac{\pi}{a}} = \frac{1}{2\sqrt{\pi}} \frac{1}{\sqrt{a}}
 \end{aligned}$$

$$1 \leq x \leq 2 \Rightarrow \pi \leq \pi x^2 \leq 4\pi$$

$\therefore$  shown



8(ii)

$$\begin{aligned}
 E(\pi X^2) &= \pi E(X^2) \\
 &= \pi \int_1^2 x^2 \, dx \\
 &= \pi \left[ \frac{x^3}{3} \right]_1^2 \\
 &= \frac{7\pi}{3}
 \end{aligned}$$

9	$X \sim \text{Po}(\lambda)$ $P(Y = r) = \sum_{m=0}^{\infty} [P(X = 0)]^m P(X = r)$ $= P(X = r) \sum_{m=0}^{\infty} (e^{-\lambda})^m$ $= P(X = r) \frac{1}{1 - e^{-\lambda}}$ $E(Y) = \frac{1}{1 - e^{-\lambda}} \sum_{r=1}^{\infty} r P(X = r)$ $= \frac{1}{1 - e^{-\lambda}} \sum_{r=0}^{\infty} r P(X = r)$ $= \frac{\lambda}{1 - e^{-\lambda}}$ <p><math>\lambda</math> is small.</p> $\frac{\lambda}{1 - e^{-\lambda}} \approx \frac{\lambda}{1 - \left(1 - \lambda + \frac{\lambda^2}{2}\right)}$ $= \lambda \left( \lambda - \frac{\lambda^2}{2} \right)^{-1}$ $= \lambda \lambda^{-1} \left( 1 - \frac{\lambda}{2} \right)^{-1}$ $\approx 1 + \frac{\lambda}{2}$
9(i)	$R \sim \text{Geom}(1 - P(X = 0))$ $R \sim \text{Geom}\left(\frac{1}{2}\right)$
9(ii)	$E(R_1 + R_2) = 2E(R)$ $= 2(2) = 4$

10(a)	<p>Let <math>X</math> and <math>Y</math> be a random candidate's marks in P1 and P2 respectively.</p> <p><math>H_0</math> : median of <math>(X - Y) = 0</math></p> <p><math>H_1</math> : median of <math>(X - Y) &lt; 0</math></p> <p>Test at 5% sig level.</p> <p>Sign of <math>(X - Y)</math> : - - - - + - - -</p> <p>1 '+' , 7 '-'</p> <p>Let <math>A</math> be the no. of positive signs out of 8.</p> <p>Under <math>H_0</math>, <math>A \sim B\left(8, \frac{1}{2}\right)</math></p> <p><math>P(A \leq 1) = 0.035156 &lt; 0.05</math></p> <p>Hence we reject <math>H_0</math> as there is suff evidence at 5 % sig level that P2 is easier than P1.</p>																																				
10(b)	<table><tr><td>Candidate</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td></tr><tr><td><math>d_i = x_i - y_i</math></td><td>-28</td><td>-4</td><td>-40</td><td>-28</td><td>36</td><td>-24</td><td>-8</td><td>-16</td></tr><tr><td>Rank of <math> d_i </math></td><td>5.5</td><td>1</td><td>8</td><td>5.5</td><td>7</td><td>4</td><td>2</td><td>3</td></tr><tr><td>Sign</td><td>-</td><td>-</td><td>-</td><td>-</td><td>+</td><td>-</td><td>-</td><td>-</td></tr></table> <p><math>T_+ = 7</math></p> <p><math>T_- = 29</math></p> <p><math>T_{calc} = \min(7, 29) = 7 &gt; 5</math></p> <p>Hence do not reject <math>H_0</math> as there is insuff evidence at 5% sig level that P2 is easier than P1.</p>	Candidate	1	2	3	4	5	6	7	8	$d_i = x_i - y_i$	-28	-4	-40	-28	36	-24	-8	-16	Rank of $ d_i $	5.5	1	8	5.5	7	4	2	3	Sign	-	-	-	-	+	-	-	-
Candidate	1	2	3	4	5	6	7	8																													
$d_i = x_i - y_i$	-28	-4	-40	-28	36	-24	-8	-16																													
Rank of $ d_i $	5.5	1	8	5.5	7	4	2	3																													
Sign	-	-	-	-	+	-	-	-																													
	<p>Wilcoxon test is more appropriate. It takes into account both the magnitude and sign of the differences in marks for each candidate.</p>																																				