

TEMASEK JUNIOR COLLEGE, SINGAPORE  
JC 2  
Preliminary Examination 2017  
Higher 2



**FURTHER  
MATHEMATICS  
Paper 1**

**9649/01**

**30 August 2017**

Additional Materials:     Answer paper  
                                     List of Formulae (MF26)

**3 hours**

---

**READ THESE INSTRUCTIONS FIRST**

Write your Civics group and name on all the work that you hand in.

Write in dark blue or black pen on both sides of the paper.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

You are expected to use a graphic calculator.

Unsupported answers from a graphic calculator are allowed unless a question specifically states otherwise.

Where unsupported answers from a graphic calculator are not allowed in a question, you are required to present the mathematical steps using mathematical notations and not calculator commands.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

This document consists of 7 printed pages and 1 blank page.

- 1** The harmonic numbers  $H_k$ ,  $k = 1, 2, 3, \dots$ , are defined by

$$H_k = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{k}.$$

Prove by mathematical induction that

$$H_{2^n} \geq 1 + \frac{n}{2}, \text{ where } n \text{ is a non-negative integer.} \quad [6]$$

- 2** Show, by means of the substitution  $u = y^{1-n}$ , that the Bernoulli's differential equation of the form

$$\frac{dy}{dx} + f(x)y = g(x)y^n, \text{ where } n \text{ is a non-zero integer and } n \neq 1$$

can be reduced to the form  $\frac{du}{dx} + P(x)u = Q(x)$ . [2]

A cardiac pacemaker is designed to provide electrical impulses  $I$  amps such that as time  $t$  increases,  $I$  oscillates with a fixed amplitude of one amp. It is proposed that the following differential equation

$$\frac{dI}{dt} + (\tan t)I = (I \sin t)^2$$

can be used to describe how  $I$  changes with  $t$ .

By using a substitution of the form  $u = I^{1-n}$ , find  $I$  in terms of  $t$ . [5]

State one limitation of this model. [1]

- 3** (a) A computer system considers a string of digits a valid codeword if it contains an even number of 0 digits. For example, 123040826 and 14947 are valid codewords, whereas 9038040 is not. Let  $a_n$  be the number of valid  $n$ -digit codewords.


(i) Find the values of  $a_1$  and  $a_2$ . [2]

(ii) By considering the number of valid and invalid  $(n-1)$ -digit codewords, find a recurrence relation for  $a_n$ . [2]

- (b) Solve the recurrence relation  $b_n = 4b_{n-1} - 4b_{n-2}$  for  $n \geq 2$  with initial conditions

$$b_0 = 6 \text{ and } b_1 = \sqrt{5}. \quad [6]$$

4



The diagram shows the vertical cross-section of a specially-designed sound reflector hanging from the ceiling of a room. The arc  $ACB$  forms part of an ellipse with  $F$  as one of the foci.  $CF$  is a vertical line of symmetry and  $AFB$  is a horizontal straight line 3 m above the ground.

A sound transmitter is positioned at  $F$  and a sound receiver is to be placed at the other focus  $X$  of the ellipse so as to receive the maximum intensity of the sound from the transmitter.

- (i) Given that  $CF = 12$  cm and  $AB = 40$  cm, determine the position of  $X$ . [3]
- (ii) Find the polar equation of the ellipse if  $F$  is positioned at the pole and  $FB$  is in the direction of the line  $\theta = 0$ . [4]

5 Let  $V$  and  $W$  be subspaces of  $\mathbb{R}^n$ . Show that  $V \cap W$  is also a subspace of  $\mathbb{R}^n$ . [3]

The transformation  $T_1: \mathbb{R}^4 \rightarrow \mathbb{R}^3$  and  $T_2: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  are represented by the matrices

$$\mathbf{M} = \begin{pmatrix} 1 & 0 & -1 & 2 \\ 2 & 1 & -1 & 3 \\ 1 & 2 & 1 & 0 \end{pmatrix} \text{ and } \mathbf{N} = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} \text{ respectively. The range space of } T_1 \text{ is}$$

denoted by  $A$  and the range space of  $T_2$  is denoted by  $B$ .

- (i) Find the rank of  $\mathbf{M}$  and find a basis for  $A$ . [3]
- (ii) Write down a basis for  $B$ . [1]
- (iii) Find a basis for  $A \cap B$ . [3]
- (iv) Determine whether  $A \cup B$  is a subspace of  $\mathbb{R}^3$ . Justify your answer. [4]

[Turn over

- 6 (a) (i)** Solve the equation

$$z^5 - 32i = 0,$$

giving the roots in the form  $re^{i\alpha}$ , where  $r > 0$  and  $-\pi < \alpha \leq \pi$ .

Draw and represent these roots in an Argand diagram. [4]

- (ii)** Given that  $w = \frac{1}{2}ze^{-i\frac{\pi}{10}}$ , describe geometrically how the point representing  $w$

can be obtained from  $z$ . [1]

Find, in simplified form, an equation for which the values of  $w$  are the roots.

[1]

- (b)** The complex number  $z$  satisfies the relations  $|z - 2i| \leq 1$  and  $\arg(z + 1) \leq \frac{\pi}{4}$ .


- (i)** Illustrate the locus of the point representing  $z$  on an Argand diagram. [2]

- (ii)** Label on your Argand diagram the point  $P$  for which  $\arg(z)$  has the least value. Find this least value and hence find the coordinates of  $P$ . [5]

- 7 The figure below shows the cross-section of a proposed underground tunnel. The equation of the curved section is

$$y = \begin{cases} x - x \ln x, & x > 0 \\ 0 & , x = 0 \end{cases}$$

where  $x$  and  $y$  are in metres.



The engineers need to estimate the area of the cross section  $A \text{ m}^2$  so that they can work out the volume of the soil that will need to be removed when a straight tunnel is built.

- (i) Engineer Yang uses the trapezium rule to estimate  $A$  with 6 strips. Find his estimate correct to 3 decimal places. [2]
- (ii) Engineer Tan and Engineer Liu estimate  $A$  with 8 strips and 16 strips respectively using trapezium rule. The estimates obtained are 1.799 and 1.833. Without any calculation, identify the estimate that is obtained by engineer Tan and state your reasons clearly. [1]

In order to account for the error when using trapezium rule for the estimation of  $A$ , it is proposed to refine the formula using

$$A \approx A(n) + \frac{a}{n^2}$$

where  $A(n)$  is the estimation based on  $n$  strips and  $a$  is a positive constant.

Find a better estimation of  $A$  using the values of  $A(8)$  and  $A(16)$ . [2]

- (iii) Engineer Shin uses Simpson's rule with 7 ordinates to estimate  $A$ . Find his estimate correct to 3 decimal places. [2]

Without further calculation, explain whether Engineer Yang or Engineer Shin gives a better approximation to  $A$ . [1]

Another tunnel with the same cross-section is built by rotating the cross-section by  $2\pi$  radians about the  $y$ -axis.

- (iv) Find the exact volume of soil that need to be removed to form the loop. [4]

**[Turn over**

- 8** A study is done to find out how the population of a plant species changes under different environmental conditions. Let  $P$  denote the size of the population  $t$  months after the study began. It is given that

$$\frac{dP}{dt} = kP(N - P)$$

where  $k$  is a positive constant and  $N$  is the maximum sustainable population of the plant species in the region.

Using an appropriate sketch of  $P$  against  $t$  for each of the following on a single diagram, explain without solving the differential equation, how  $P$  changes with  $t$  when

- (i)  $p_0 > N$                       (ii)  $0 < p_0 < N$

where  $P = p_0$  is the initial population size. [4]

It is given that  $p_0 = 20$  and  $k = 0.01$ .

- (a) When  $N = 1500$ , Euler's method of step size  $\Delta t = 0.5$  month is used to estimate the population sizes. The following table of result is obtained where  $P = p_n$  and  $f(t_n, p_n) = kp_n(N - p_n)$  after  $t_n$  months.

$n$	$t_n$	$p_n$	$f(t_n, p_n)$
0	0	20	266
1	0.5	168	1831
2	1	1287	2742


Explain, with the aid of an appropriate sketch, whether the population size estimated using Euler's method at the end of one month is an over or under-estimation. [3]

Explain whether Euler's Method can be used to estimate the population size at the end of the second month. [2]

- (b) Due to the presence of locust during different seasons,  $N$  varies with  $t$  and is given by

$$N = 1500 \left[ 1 - \frac{1}{10} \cos\left(\frac{1}{4}t\right) \right]$$

Use Improved Euler's Method once to estimate the population size at the end of one month. [3]



The polar equation of the curve  $C_1$  is

$$r = 3 + 2 \cos \theta, \quad 0 \leq \theta \leq 2\pi.$$

The circle  $C_2$  with centre at the pole  $O$  and radius 2 units intersects  $C_1$  at the points  $P$  and  $Q$ , as shown in the diagram.

- (a) Find the polar coordinates of  $P$  and  $Q$ . [3]
- (b) The straight line  $PO$  is extended to intersect the curve again at the point  $A$ .
  - (i) Find the polar coordinates of  $A$ . [2]
  - (ii) Find the exact length of  $AQ$ . [3]
  - (iii) Hence, or otherwise, show that the line  $AQ$  is a tangent to the circle  $C_2$ . [2]
- (c) The shaded region  $R$  lies inside  $C_2$  but outside  $C_1$ . Show that the area of  $R$  can be expressed in the form  $\frac{1}{6}(a\sqrt{3} + b\pi)$ , where  $a$  and  $b$  are integers to be determined. [8]